

## Goat Guideline for Anthelmintic Dosages (internal parasite dewormers) July 2006

***\*Important --- Please read notes below before using this chart\****

Animal Weight		Oral dosing.					Subcutaneous injection	
		Valbazen Albendazole <sup>1</sup>	SafeGuard Fenbendazole <sup>2</sup>	Ivomec Ivermectin <sup>3</sup>	Levasole Levamisole <sup>4</sup>	Cyductin Pour on Moxidectin <sup>5</sup>	Cyductin Drench Moxidectin <sup>6</sup>	Cyductin *Injectable* Moxidectin <sup>7</sup>
lbs	kg	20 mg/kg 2 ml/ 25 lb	10 mg/kg 1.1 ml/ 25 lb	0.4 mg/kg 6 ml/ 25 lb	12 mg/kg 3 ml/ 25 lb	0.5 mg/kg 1.1 ml/25 lb	0.3 mg/kg 3.4 ml/25 lb	0.2 mg/kg 1 ml/ 110 lb
20	9.1	1.6	0.9	4.8	2.4	0.9	2.7	0.2
25	11.4	2.0	1.1	6.0	3.0	1.1	3.4	0.2
30	13.6	2.4	1.4	7.2	3.6	1.4	4.1	0.3
35	15.9	2.8	1.6	8.4	4.2	1.6	4.8	0.3
40	18.2	3.2	1.8	9.6	4.8	1.8	5.4	0.4
45	20.5	3.6	2.1	10.8	5.4	2.1	6.1	0.4
50	22.7	4.0	2.3	12.0	6.0	2.3	6.8	0.5
55	25.0	4.4	2.5	13.2	6.6	2.5	7.5	0.5
60	27.3	4.8	2.7	14.4	7.2	2.7	8.2	0.5
65	29.5	5.2	3.0	15.6	7.8	3.0	8.8	0.6
70	31.8	5.6	3.2	16.8	8.4	3.2	9.5	0.6
75	34.1	6.0	3.4	18.0	9.0	3.4	10.2	0.7
80	36.4	6.4	3.6	19.2	9.6	3.6	10.9	0.7
85	38.6	6.8	3.9	20.4	10.2	3.9	11.6	0.8
90	40.9	7.2	4.1	21.6	10.8	4.1	12.2	0.8
95	43.2	7.6	4.3	22.8	11.4	4.3	12.9	0.9
100	45.5	8.0	4.6	24.0	12.0	4.6	13.6	0.9
105	47.7	8.4	4.8	25.2	12.6	4.8	14.3	1.0
110	50.0	8.8	5.0	26.4	13.2	5.0	15.0	1.0
115	52.3	9.2	5.2	27.6	13.8	5.2	15.6	1.0
120	54.5	9.6	5.5	28.8	14.4	5.5	16.3	1.1
125	56.8	10.0	5.7	30.0	15.0	5.7	17.0	1.1
130	59.1	10.4	5.9	31.2	15.6	5.9	17.7	1.2
140	63.6	11.2	6.4	33.6	16.8	6.4	19.0	1.3
150	68.2	12.0	6.8	36.0	18.0	6.8	20.4	1.4

<sup>1</sup>**Valbazen** Suspension (11.36 % or 113.6 mg/ml): *Do NOT use in pregnant does in the first trimester of pregnancy.* Meat withdrawal time is 9 days and 7 days for milk (FARAD).

<sup>2</sup>**Safe-Guard/ Panacur** Suspension (10% or 100 mg/ml): Approved in goats at 5 mg/kg with meat withdrawal time of 6 days and no withdrawal period for milk. Although the label dose in goats is 5 mg/kg, it is generally recognized that 10 mg/kg dosage is required for good efficacy. At 10 mg/kg dosage, meat withdrawal is 16 days and 4 days for milk (FARAD).

<sup>3</sup>**Ivomec Sheep Drench** (0.08% or 0.8 mg/ml): Protect from light. Coughing may occur during and following drenching. Meat withdrawal time is 14 days (FARAD).

<sup>4</sup>**Levasole Soluble Drench Powder (Sheep)**: Oral solution ONLY. To prepare use 1 packet (13 gm/11.7 gm active ingredient) dissolved in 262 ml [8.9 oz.] water (44.7 mg/ml) {or 52 gram packet dissolved in 1048 ml water [35.4 oz.].} NOTE: This is different dilution from the label directions for administration. Meat withdrawal time is 4 days (FARAD).

<sup>5</sup>**Cydectin Pour on for cattle** (0.5% or 5 mg/ml): Meat withdrawal time is 23 days. *Not for use in lactating dairy goats.*

<sup>6</sup>**Cydectin Drench for sheep** (.1% or 1 mg/ml): Meat withdrawal time is 14 days. *Not for use in lactating dairy goats.*

<sup>7</sup>**Cydectin Injectable for cattle** (1% or 10mg/ml): GIVE SQ. Meat withdrawal time is 30 days. *Not for use in lactating dairy goats.*

## **NOTE for Guideline for Anthelmintic Dosages in Goats**

The attached chart was developed by Ray M. Kaplan, DVM, PhD (University of Georgia) and modified by Patty Scharko DVM, MPH (University of Kentucky.) and Lionel Dawson DVM, PhD. (Oklahoma State University). It is provided as a possible guideline for anthelmintic (deworming) dosages for goats. Producers should consult their veterinarian for advice on their specific management situation for determining dosages for their herd. *With the exception of fenbendazole administered at the 5 mg/kg dose, these drugs are not approved by the Food and Drug Administration (FDA) for use in goats, and when used in goats are considered extra-label use (fenbendazole at the recommended dose rate of 10 mg/kg is considered extra-label usage). The FDA regards extra-label use of drugs as an exclusive privilege of the veterinary profession and is only permitted when a bona fide veterinarian-client-patient relationship exists and an appropriate medical diagnosis has been made. The chart is intended to serve as guideline for improving accuracy when dosing goats with an anthelmintic, but these drugs should be used in goats only when appropriate veterinary advice has been received.*

Drug resistance in parasites of goats is extremely common. The effectiveness of an anthelmintic should always be tested before being used by performing a FECRT (Fecal Egg Count Reduction test) or larval development (DrenchRite) assay if available.

*\*\* The current recommendation is to use the Cydectin cattle **injectable** formulation and **NOT** the **pour-on** formulation (orally) or the sheep oral drench. When administered by subcutaneous injection, moxidectin provides improved drug levels as compared to when administered orally.*

---

*Information contained in this document is part of a web-based training and certification program for meat goat producers (<http://www2.luresext.edu/goats/training/ga.html>) that was developed with funding received by Langston University from USDA/FSIS/OPHS project #FSIS-C-10-2004 entitled "Development of a Web-based Training and Certification Program for Meat Goat Producers."*

Collaborating institutions/organizations include Alcorn State University, American Boer Goat Association, American Kiko Goat Association, American Meat Goat Association, Florida A&M University, Fort Valley State University, Kentucky State University, Langston University, Prairie View A&M University, Southern University, Tennessee Goat Producers Association, Tennessee State University, Tuskegee University, United States Boer Goat Association, University of Arkansas Pine Bluff, and Virginia State University.